

MANAGEMENT STRATEGIES



Dow AgroSciences

HOW TO TREAT

1. TREATING ISOLATED PATCHES AND LIMITED INFESTATIONS UNDER EUCALYPT TREES

Use Starane™ Advanced Herbicide to control mother-of-millions present under eucalypt trees. Starane Advanced has no soil residue and will not affect the eucalypt trees.

2. TREATING EXTENSIVE INFESTATIONS ON ARABLE LAND

Use Grazon™ Extra Herbicide as a foliar spray to control mother-of-millions in open areas. Grazon Extra will suppress new seedlings. Encourage pasture growth to help with competition.

3. TREATING EXTENSIVE INFESTATIONS ON HILL COUNTRY

Containment of mother-of-millions to a particular site is the first strategy. The next step is to reduce the infestation by using foliar application on smaller patches first, then moving on to larger patches. It may be beneficial to fence off areas so cattle do not graze these areas during flowering (winter months).



FOR BEST RESULTS

Use Grazon Extra on mother-of-millions when flowering at 500 mL/100 L of water rate. Add a 100% concentrate non-ionic surfactant (e.g. BS1000®) at 100 mL/100 L of water. When spraying, treat the soil around the mother-of-millions as Grazon Extra has some residual activity in the soil and will control new seedlings up to six months after application.

Use Starane Advanced on seedlings and young plants before flowering at 360 mL/100 L of water with a surfactant. Starane Advanced has no soil residual activity and can be safely used among eucalypt trees.

Application technique: use spot spray technique to thoroughly wet leaves and stems to the point of run-off using a handgun with a D6 nozzle at an operating pressure of 500–800 kPa. A narrow spray jet penetrates the clump and wets stems while leaves are coated using a wider cone. Always spray in a systematic pattern ensuring thorough coverage of each clump before moving to the next.

TREATMENT TIPS

Poisonous plants may become more palatable after spraying and stock should be kept away from these plants until they die down.

Always use a 100% non-ionic surfactant at the rate of 100 mL/100 L of water.

MOTHER-OF-MILLIONS

(*Bryophyllum* spp.)



MOTHER-OF-MILLIONS (*Bryophyllum* spp.)

DESCRIPTION

Mother-of-millions are erect, smooth, fleshy succulent plants growing to 1 m tall or more. All species form tall flower spikes in winter with clusters of bell-shaped flowers. Each species has a distinctive leaf-shape but all produce small plantlets along the edges of the leaves. These plantlets drop readily, develop roots and establish quickly to form a new colony. Of the three species naturalised in Queensland, two produce tall flower spikes in winter and have orange-red bell-shaped flowers that form in clusters at the top of a single stem. The third type, resurrection plant or green mother-of-millions, produces yellowish-green flowers, usually with a pink tinge that grow in clusters along the upper portion of the stem.

Plant material is poisonous to livestock, flowers are especially toxic if ingested by grazing animals. Cases of cattle poisoning have been reported mainly during drought years when grass was very dry and other green feed unavailable. The main symptoms reported in cattle are scouring or straining, sometimes with evidence of abdominal pain and the faeces blood stained or mucus-covered, with many affected animals dying after two to three days.



HERBICIDES FOR CONTROL OF MOTHER-OF-MILLIONS

PRODUCT#	METHOD OF APPLICATION	RATE†	STATE
Grazon™ Extra Herbicide	Foliar spray	500 mL/100 L of water + surfactant	All
Starane™ Advanced Herbicide	Foliar spray	360 mL/100 L of water + surfactant	All
Statesman™ 720 Herbicide	Foliar spray	350 mL/100 L of water + surfactant	ACT, NSW

Note: Preferred product option in bold. † Consult label for details of growth stage and use, especially where range of rates is given.

TIMING

Mother-of-millions can be controlled with herbicides at any time of the year but infestations are easiest to see in winter when the plants are in flower. Treating infestations at this time of year also has the benefit of preventing new seeds from developing.

PRODUCT	Sep	Oct	Nov	Dec	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	KEY
Grazon Extra	⚙️	⚙️	⚙️	⚙️	⚙️	⚙️	⚙️	⚙️	✓	✓	✓	✓	Best time to spray/treat ✓
Starane Advanced	⚙️	⚙️	⚙️	⚙️	⚙️	✓	✓	✓	⚙️	⚙️	⚙️	⚙️	Can spray/treat if conditions are suitable ⚙️



Confidence in a drum

FOR MORE INFORMATION CALL
TOLL FREE 1800 700 096 OR VISIT
www.dowagrosciences.com.au